

Anti-stressant

Resoil reanimator

Universal natural product derived from leonardite with a high content of fulvic acids, which is used both for seed treatment and for foliar feeding.

Contributes to the active division of cells, formation of generative organs, stimulation of immunity and photosynthesis, as well as increasing plant resistance to stress of various natures. Fulvic acids, which are the part of the product, are a natural activator of growth processes.

Applied for adjusting the nutrition, stimulation and development of plant growth, activation of protective functions in the plants and increase germination and energy of seeds.

Product features

Indicator name	Units of measure	Value*	Note
Density	kg/l	1,05-1,37	
Boiling point	°C	+ 97	
Freezing point	°C	-4 -8	
Storage temperature	°C	+ 10 +45	
Temperature of loss of the properties	°C	Less than + 5 over + 50	Avoid long-term exposure to direct sunlight
Form		Liquid	
Color		From light brown with a pink tinge to dark brown	
Odor		none	
pH	pH	5,00-5,25	
Humic acids	%	0,03-0,05	
Fulvic acids	%	0,65-0,8	
Ash content	g/l	4,55	
Dry residue	g/l	5,99	
Humic acids	g/l	0,17	
Fulvic acids	g/l	0,7	
N	%	0,2-0,25	
K	mg/l	95,0	
Si	mg/l	718,6	
Ca	mg/l	1637,8	
S	mg/l	979,1	
Mn	mg/l	5,2	
Fe	mg/l	645,0	
Cu	mg/l	3,3	
Zn	mg/l	5,6	
Organic matter	g/l	5,7-5,9	

* The value of the indicators depending on the raw material may differ $\pm 10\%$

Benefits

1. 100% soluble.
2. High biological activity.
3. Environmental-friendly.
4. Rapid removal of plants from the stress - the restoration of growth process.
5. Increases the absorption of nutrients from soil and fertilizers.
6. Provides active development of the secondary root system.
7. Stimulates the rapid onset of vegetative mass.
8. Strengthens the immune system of the plants.
9. Seeds treated before sowing significantly increase the viability and energy of growth.
10. Used with different types of sprayers and drip systems.
11. Increases the resistance of plants against stress from the frost and drought (especially with sudden temperature changes) and from plant protection products.
12. Stimulates tillering and branching.
13. Intensifies metabolic processes in the plant cell (especially division of cells), increases the intensity of photosynthesis.

Application rate for the cultures:

- Gardens 2.0-7.0 l/ha (up to 2.5% in the solution with foliar application and up to 5.0% with drip irrigation).
- Greenhouse (ferrigation) 1.0-7.0 l/ha (up to 5.0% in the solution).
- Field crops:
 1. Seed treatment 1.0-5.0 l/t (depending on the surface area of the seed. For example: Rapeseed 5.0 l/t, and soybeans 1.0 l/t).
 2. Application on vegetative parts of plants:
 - Cereals 3.0-5.0 l/ha (up to 2.5% in the solution).
 - Leguminous 1.0-3.0 l/ha (up to 2.5% in the solution).
 - Vegetables 2.0-7.0 l/ha (up to 2.5% in the solution for foliar application and up to 5.0% for drip irrigation).
 - Other agricultural crops 1.0-5.0 l/ha (up to 2.5% in the solution).